



CATEGORY:

CLEARED

ADDRESS
CONTACT IF FOUND:

BO 41497

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

09/647596

INTERNATIONAL APPLICATION NO.
PCT/NL98/00186INTERNATIONAL FILING DATE
3 April 1998PRIORITY DATE CLAIMED
--

TITLE OF INVENTION

METHOD OF INTERSTRAIN DIFFERENTIATION OF BACTERIA

APPLICANT(S) FOR DO/EO/US Johannes Dirk Anthonie VAN EMBDEN,
Leendert Marinus SCHOOLS and Rudolph JANSEN

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☒ This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).
4. ☒ A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
 - a. ☒ is transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ has been transmitted by the International Bureau.
 - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ A translation of the International Application into English (35 U.S.C. 371(c)(2)).
7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
 - a. ☐ are transmitted herewith (required only if not transmitted by the International Bureau).
 - b. ☐ have been transmitted by the International Bureau.
 - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
 - d. ☐ have not been made and will not be made.
8. ☐ A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ A translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

Items 11. to 16. below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A **FIRST** preliminary amendment.
☐ A **SECOND** or **SUBSEQUENT** preliminary amendment
14. ☐ A substitute specification.
15. ☐ A change of power of attorney and/or address letter.
16. ☒ Other items or information:

International Preliminary Examination Report.

Application Data Sheet.

Search Report.

U.S. APPLICATION NO. 09/647596

INTERNATIONAL APPLICATION NO.
PCT/NL98/00186ATTORNEY'S DOCKET NUMBER
BO 4149717. ☒ The following fees are submitted:

BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)):

Neither international preliminary examination fee (37 CFR 1.482)
nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO
and International Search Report not prepared by the EPO or JPO \$ 1000.International preliminary examination fee (37 CFR 1.482) not paid to
USPTO but International Search Report prepared by the EPO or JPO 860.International preliminary examination fee (37 CFR 1.482) not paid to USPTO
but international search fee (37 CFR 1.445(a)(2)) paid to USPTO 710.International preliminary examination fee (37 CFR 1.482) paid to USPTO
but all claims did not satisfy provisions of PCT Article 33(1)-(4) 690.International preliminary examination fee (37 CFR 1.482) paid to USPTO
and all claims satisfied provisions of PCT Article 33(1)-(4) 100.

ENTER APPROPRIATE BASIC FEE AMOUNT =

CALCULATIONS PTO USE ONLY

\$ 860

Surcharge of \$130.00 for furnishing the oath or declaration later than ☐ 20 ☒ 30
months from the earliest claimed priority date (37 CFR 1.492(e)).

\$ 130

CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE
Total claims	25 - 20 =	5	x \$ 18.
Independent claims	4 - 3 =	1	x .80.
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ 270.

\$

\$ 90

\$ 80

\$

TOTAL OF ABOVE CALCULATIONS =

\$ 1,160

Reduction of 1/2 for small entity

\$

SUBTOTAL =

\$ 1,160

Processing fee of \$130.00 for furnishing the English translation later than ☐ 20 ☐ 30
months from the earliest claimed priority date (37 CFR 1.492(f)).

\$

TOTAL NATIONAL FEE =

\$ 1,160

Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be
accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property +

\$

TOTAL FEES ENCLOSED =

\$ 1,160

Amount to be
refunded:

\$

charged:

\$

a. ☒ A check in the amount of \$ 1,160 to cover the above fees is enclosed.b. ☐ Please charge my Deposit Account No. _____ in the amount of \$ _____ to cover the above fees.
A duplicate copy of this sheet is enclosed.c. ☒ The Commissioner is hereby authorized to charge any additional fees which may be required by
37 CFR 1.16 and 1.17, or credit any overpayment to Deposit Account No. 25-0120. A duplicate
copy of this sheet is enclosed.NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR
1.137 (a) or (b)) must be filed and granted to restore the application to pending status.

October 3, 2000

SEND ALL CORRESPONDENCE TO:

Young & Thompson
745 South 23rd Street
2nd Floor
Arlington, VA 22202
(703) 521-2297

CUSTOMER NO. 000466

SIGNATURE

Benoit Castel
NAME35,041
REGISTRATION NUMBER

09/647596

532 Rec'd PCT/PTC 03 OCT 2000

PATENTS

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of

Johannes D.A. VAN EMBDEN et al.

Serial No. (unknown)

Filed herewith

METHOD INTERSTRAIN DIFFERENTIA-
TION OF BACTERIA

PRELIMINARY AMENDMENT

Commissioner for Patents

Washington, D.C. 20231

Sir:

Prior to calculation of the filing fee, please amend
the above-identified application as follows:

IN THE DRAWINGS:

Please add Figure 6 as filed in the Article 34
amendment of June 2, 2000. The sheet containing Figure 6 is
attached hereto and marked "AMENDED SHEET".

IN THE CLAIMS:

Please substitute Claims 1-25 as originally filed,
which appear on pages 22-27, with Claims 1-25 as filed in the
Article 34 amendment of June 15, 2000. The pages containing
Claims 1-25 are marked "AMENDED SHEET" and are attached
hereto. Following the insertion of Claims 1-25, please amend
these claims as follows:

Claim 3, line 1, cancel "or 2".

*6.6 AM 34
SUB CLAIM AMENDED
by PRAHAR - PCT*

Johannes D.A. VAN EMBDEN et al.

Claim 4, line 1, change "any of the preceding claims" to --claim 1--.

Claim 5, line 1, change "any of the preceding claims" to --claim 1--.

Claim 6, line 1, change "any of the preceding claims" to --claim 1--.

Claim 7, line 1, change "any of the preceding claims" to --claim 1--.

Claim 8, line 1, change "any of the preceding claims" to --claim 1--.

Claim 10, line 1, change "any of the preceding claims" to --claim 1--.

Claim 11, line 1, change "any of the preceding claims" to --claim 1--.

Claim 14, line 1, cancel "or 13".

Claim 15, line 1, change "any of claims 12-14" to --claim 12--.

Claim 16, line 1, change "any of claims 12-15" to --claim 12--.

Claim 17, line 3, change "any of claims 12-16" to --claim 12--.

Claim 19, line 1, cancel "or 18".

Claim 21, line 1, change "claim 21" to --claim 20--.

Amend Claim 24 as follows:

--24. (amended) Kit for carrying out a method [according to any of claims 1-19] of in vitro amplification of nucleic acid using amplification primers, comprising a primer pair according to [any of claims 20-23] claim 20 and an

Johannes D.A. VAN EMBDEN et al.

oligonucleotide probe or a carrier, said carrier comprising at least 1 oligonucleotide probe specific for a spacer region of a bacterium to be determined said bacterium not belonging to M. tuberculosis complex, said oligonucleotide probe being an oligonucleotide probe comprising between 12 to 40 nucleotides, said probe being sufficiently homologous to any of the spacer sequences or to fragments or derivatives of such spacer sequences to hybridise to such a spacer sequence, said oligonucleotide probe comprising at least ten consecutive nucleotides homologous to such a spacer sequence and/or exhibiting at least 60% homology, with the corresponding part of the spacer sequence.--

R E M A R K S

The above changes in the drawings and claims merely place the national phase application in the same condition as it was during Chapter II of the international phase, with the multiple dependencies in the claims being removed.

Respectfully submitted,

YOUNG & THOMPSON

By Benoit Castel
Benoit Castel
Attorney for Applicants
Registration No. 35,041
745 South 23rd Street
Arlington, VA 22202
Telephone: 703/521-2297

October 3, 2000

02-06-2000

NL 009800186

1

Claims

1. A method of in vitro amplification of nucleic acid using amplification primers in a manner known per se, in amplification reactions such as PCR, LCR or NASBA, wherein
5 a pair of primers is used comprising oligonucleotide sequences sufficiently complementary to a part of the Direct Repeat sequence of a bacterium other than a microorganism belonging to the Mycobacterium tuberculosis complex of microorganisms for hybridisation to a Direct Repeat to occur and subsequently elongation of the hybridised primer to take place, said primers being such that elongation
10 in the amplification reaction occurs for one primer in the 5' Direction and for the other primer in the 3' Direction, wherein the Direct Repeat is a sequence with a length between 20-50 base pairs which occurs 5-60 times in a region of the bacterial genome, whereby the Direct Repeat sequences are separated by spacer sequences with a length of between 20-50 nucleotides, said spacer sequences being non repetitive.
- 15 2. A method according to claim 1 wherein the Direct Repeat sequence is obtainable from screening a genomic bacterial nucleic acid sequence using the programme Patscan wherein the Direct Repeat is designated p1 with a length between 20-50 basepairs then p1 is sought 20-50 basepairs downstream of p1 as the pattern

02-06-2000

NL 009800186

2

p1=(20..50)(20..50)p1(20..50)p1 or a variant thereof wherein the ranges of the nucleotide lengths are shorter and wherein the frequency of occurrence of the Direct Repeat can vary between 5 and 60.

- 5 3. A method according to claim 1 or 2 wherein the Direct Repeat has a length between 30-40 base pairs and the spacer has a length of 35-45 base pairs.
4. A method according to any of the preceding claims wherein the Direct Repeat has a terminus exhibiting at least 3 out of 5 nucleotides identical with the
10 sequence GAAAC, preferably 4, said termini being selected from GAAAC, GAAXXC, GAACTC, GXAAC, GCAAC, GAAA, GAAXC, GAAGC, AAAC.
5. A method according to any of the preceding claims wherein the Direct Repeat terminates with GAACTC, ATACAC, AAAACT, TTGCAA, GGAAAC,
15 TGAAAC, TGAAGC, TGGAAA, TTAAAC, TGAAAT or TTCAAC.
6. A method according to any of the preceding claims wherein the Direct Repeat has stretches of 3-4 identical bases.
- 20 7. A method according to any of the preceding claims wherein the Direct Repeat has a sequence such that it is not prone to loop formation or any other obvious secondary structure.
8. A method according to any of the preceding claims wherein the bacterium is
25 a pathogenic bacterium selected from the group of Gram negative aerobic/microaerophilic rods and cocci and facultatively anaerobic gram-negative rods.
9. A method according to claim 8 wherein the bacterium is a pathogenic bacterium selected from the genera Escherichia, Shigella, Salmonella, Klebsiella,
30 Enterobacter, Yersinia, Serratia, Haemophilus, Vibrio, Legionella, Neisseria, Pseudomonas, Bordetella, Staphylococcus, Streptococcus and Acinetobacter.

02-06-2000

3

10. A method according to any of the preceding claims, wherein said primers have oligonucleotide sequences complementary to non overlapping parts of the Direct Repeat sequence and such that the elongation reactions from each primer can occur without hindrance of the other when both primers hybridise to the same Direct Repeat and undergo elongation.

11. A method according to any of the preceding claims, wherein one primer DRa is capable of elongation in the 5' Direction and the other primer DRb is capable of elongation in the 3' Direction and DRa is complementary to a sequence of the Direct Repeat located to the 5' side of the sequence of the Direct Repeat to which DRb is complementary.

12. A method of detection of a bacterium, said bacterium not belonging to the M tuberculosis complex of microorganisms said method comprising

15 1) amplifying nucleic acid from a sample with the method according to any of the preceding claims, followed by

2) carrying out a hybridisation test in a manner known per se, wherein the amplification product is hybridised to an oligonucleotide probe or a plurality of different oligonucleotide probes, each oligonucleotide being sufficiently

20 homologous to a part of a spacer of the Direct Region of the bacterium to be determined for hybridisation to occur to amplified product if such spacer nucleic acid was present in the sample prior to amplification, said hybridisation step optionally being carried out without prior electrophoresis or separation of the amplified product.

25 3) detecting any hybridised products in a manner known per se.

13. A method according to claim 12, wherein the hybridisation test is carried out using a number of oligonucleotide probes, said number comprising at least a number of oligonucleotides probes specific for the total spectrum of bacteria it is desired to detect.

30

02-06-2000

NL 009800186

4

14. A method according to claim 12 or 13, wherein the oligonucleotide probe is at least ten oligonucleotides long and is a sequence complementary to a sequence selected from any of the spacer sequences of the Direct Repeat region of the bacterium to be determined or is a sequence complementary to fragments or derivatives of said spacer sequences, said oligonucleotide probe being capable of hybridising to such a spacer sequence and comprising at least ten consecutive nucleotides homologous to such a spacer sequence and/or exhibiting at least 60% homology, preferably exhibiting at least 80% homology with such a spacer sequence.
- 10 15. A method according to any of claims 12-14 wherein the bacterium is a pathogenic bacterium selected from the group of Gram negative aerobic/microaerophilic rods and cocci and facultatively anaerobic Gram-negative rods.
- 15 16. A method according to any of claims 12-15 wherein the bacterium is a pathogenic bacterium selected from the genera *Escherichia*, *Shigella*, *Salmonella*, *Klebsiella*, *Enterobacter*, *Yersinia*, *Serratia*, *Haemophilus*, *Vibrio*, *Legionella*, *Neisseria*, *Pseudomonas* and *Bordetella* and the group of Gram positive bacterial genera *Staphylococcus* and *Streptococcus* as target for the differentiation method.
- 20 17. A method for differentiating the type of bacterium in a sample, said bacterium not belonging to the *M. tuberculosis* complex, said method comprising carrying out the method according to any of claims 12-16, followed by comparison of the hybridisation pattern obtained with a reference.
- 25 18. A method according to claim 17, wherein the reference is the hybridisation pattern obtained with one or more known strains of the bacterium to be determined in analogous manner.
19. A method according to claim 17 or 18 wherein the reference is a source providing a list of spacer sequences and sources thereof, such as a data bank.
- 30

02-06-2000

NL 009800186

5

20. A pair of primers wherein both primers comprise oligonucleotide sequences of at least 7 oligonucleotides and are sufficiently complementary to a part of the Direct Repeat sequence of the microorganism *E. coli* for hybridisation to occur and subsequently elongation of the hybridised primer to take place, said primers being such
5 that elongation in the amplification reaction occurs for one primer in the 5' Direction and for the other primer in the 3' Direction and wherein sufficiently complementary means said oligonucleotide sequence comprises at least seven consecutive nucleotides homologous to such a Direct Repeat sequence in particular a Sequence described in SEQ ID No. 1-18 and/or exhibits at least 60% homology, with the corresponding part of the
10 Direct Repeat sequence.

21. Primer pair according to claim 21, comprising one primer DRa capable of elongation in the 5' Direction and the other primer DRb capable of elongation in the 3' Direction with DRa being complementary to a sequence of the Direct Repeat located to
15 the 5' side of the sequence of the Direct Repeat to which DRb is complementary, the Direct Repeat being present in the Direct Region of SEQ ID No. 2.

22. A pair of primers wherein both primers comprise oligonucleotide sequences of at least 7 oligonucleotides and are sufficiently complementary to a part of the Direct
20 Repeat sequence of the microorganism *S. typhimurium* for hybridisation to occur and subsequently elongation of the hybridised primer to take place, said primers being such that elongation in the amplification reaction occurs for one primer in the 5' Direction and for the other primer in the 3' Direction and wherein sufficiently complementary means said oligonucleotide sequence comprises at least seven consecutive nucleotides
25 homologous to such a Direct Repeat sequence in particular the DEQ ID No. 2 and/or exhibits at least 60% homology, with the corresponding part of the Direct Repeat sequence.

23. Primer pair according to claim 22, comprising one primer DRa capable of
30 elongation in the 5' Direction and the other primer DRb capable of elongation in the 3' Direction with DRa being complementary to a sequence of the Direct Repeat located to the 5' side of the sequence of the Direct Repeat to which DRb is complementary, the Direct Repeat being present in the Direct Region of SEQ ID No. 8.

02-06-2000

NL 009800186

6

24. Kit for carrying out a method according to any of claims 1-19, comprising a primer pair according to any of claims 20-23 and an oligonucleotide probe or a carrier, said carrier comprising at least 1 oligonucleotide probe specific for a spacer region of a
5 bacterium to be determined said bacterium not belonging to *M. tuberculosis* complex, said oligonucleotide probe being an oligonucleotide probe comprising between 12 to 40 nucleotides, said probe being sufficiently homologous to any of the spacer sequences or to fragments or derivatives of such spacer sequences to hybridise to such a spacer sequence, said oligonucleotide probe comprising at least ten consecutive nucleotides
10 homologous to such a spacer sequence and/or exhibiting at least 60% homology, with the corresponding part of the spacer sequence.

25. Kit according to claim 24 further comprising a data carrier with required reference patterns of the bacterial strain to be determined.

A method of interstrain differentiation of bacteria.

Summary of the invention

5

The subject invention lies in the field of interstrain differentiation of bacteria. A general method has been developed with which various types of bacteria can be differentiated into separate individual strains. Thus in particular in the clinical setting this method can suitably be used to determine what strain of bacterium is present in a sample. This new method is applicable for discerning between various strains of both Gram negative and Gram positive types of bacteria.

15 Background of the invention

Previously we had disclosed a method called oligotyping for interstrain differentiation of *Mycobacterium tuberculosis* strains in W095/31569. It was stated in this document that one of the key factors in the control of tuberculosis is the rapid diagnosis of the disease and the identification of the sources of infection. *M. tuberculosis* strain typing has already proved to be extremely useful in outbreak investigations (6, 14, 31) and is being applied to a variety of epidemiologic questions in numerous laboratories. Traditionally, laboratory diagnosis is done by microscopy, culturing of the micro-organism, skin testing and X-ray imaging. Unfortunately, these methods are often not sensitive, not specific and are very time-consuming, due to the slow growth rate of *M. tuberculosis*. Therefore, new techniques like in vitro amplification of *M. tuberculosis* DNA have been developed to rapidly detect the micro-organism in clinical specimens (14). The ability to differentiate isolates of *M. tuberculosis* by DNA techniques has revolutionarized the potential to identify the sources of infection and to establish main routes of transmission and risk factors for acquiring tuberculosis by infection (1,3-10, 14, 16, 19-22, 25, 26, 27-33). The use of an effective universal typing system will allow strains from different geographic areas to be compared and the movement of individual strains to be tracked. Such data may provide important insights and identify strains with particular problems such as high infectivity, high virulence and/or multidrug resistance. Analysis of large numbers of isolates may provide answers to long-standing questions regarding the efficacy of BCG vaccination and the

40

frequency of reactivation versus reinfection.

The same problems identified for *M. tuberculosis* are inherent in differentiation of numerous other bacteria. The problems specifically arise for potentially epidemic pathogens and for bacteria that infect hospitals. A more rapid and simple typing method is required. Preferably the testing methods for various bacteria will occur in the same manner ensuring routine use for all types of bacteria for which testing is required. Preferably a test that can be carried out by non specialised personnel using little laboratory space and time is sought after.

The method disclosed in W095/31569 is based on the DNA polymorphism found at a unique chromosomal locus, the "Direct Repeat" (DR) region, which is uniquely present in *M. tuberculosis* complex bacteria. This locus was discovered by Hermans et al. (15) in *M. bovis* BCG, the strain used worldwide to vaccinate against tuberculosis. The DR region in *M. bovis* BCG consists of Directly repeated sequences of 36 base pairs, which are interspersed by non-repetitive DNA spacers, each 35 to 41 base pairs in length (15). The number of copies of the DR sequence in *M. bovis* BCG was determined to be 49. In other strains of the *M. tuberculosis* complex the number of DR elements was found to vary (15). The vast majority of the *M. tuberculosis* strains contain one or more IS6110 elements in the DR containing region of the genome.

It has been shown (12) that the genetic diversity in the DR region is generated by differences in the DR copy number, suggesting that homologous recombination between DR sequences may be a major driving force for the DR-associated DNA polymorphism (12). The high degree of DNA polymorphism within a relatively small part of the chromosome makes this region well-suited for a PCR-based fingerprinting technique.

Figure 1 depicts the structure of the DR region of *M. bovis* BCG as determined previously by Hermans et al. and Groenen et al. (12, 15). For the sake of convenience we will designate a DR plus its 3' adjacent spacer sequence as a "Direct Variant Repeat" (DVR). Thus, the DR region is composed of a discrete number of DVR's, each consisting of a constant part (DR) and a variable part (the spacer).

The method disclosed in W095/31569 is based on a unique method of in vitro amplification of DNA sequences within the DR region and the hybridisation of the amplified DNA with multiple, short synthetic oligomeric DNA sequences based on the sequences of the unique spacer DNA's within the DR region (figure 2). This differs from previous PCR methods in the use of a set of primers with both primers having multiple

priming sites as opposed to having one of the primers bind to a fixed priming site such as to a part of IS6110. Because *M. tuberculosis* complex strains differ in the presence of these spacer sequences, strains can be differentiated by the different hybridisation patterns with a set of various spacer DNA sequences.

The method consists of in vitro amplification of nucleic acid using amplification primers in a manner known per se in amplification reactions such as PCR, LCR or NASBA, wherein a pair of primers is used comprising oligonucleotide sequences sufficiently complementary to a part of the Direct Repeat sequence of a microorganism belonging to the *M. tuberculosis* complex of microorganisms for hybridisation to a Direct Repeat to occur and subsequently elongation of the hybridized primer to take place, said primer being such that elongation in the amplification reaction occurs for one primer in the 5' Direction and for the other primer in the 3' Direction. Due to the multiple presence of Direct Repeats in the microorganisms to be detected the use of such primers implies that all the spacer regions will be amplified in an efficient manner. In particular it is not necessary for extremely long sequences to be produced in order to obtain amplification of spacers located at a distance from the primer. With the instant selection of the primer pairs a heterogenous product is obtained comprising fragments all comprising spacer region nucleic acid. Subsequently the detection of the amplified product can occur simply by using an oligonucleotide probe directed at one or more of the spacer regions one wishes to detect. In order to avoid hindrance in the amplification reactions the primers can have oligonucleotide sequences complementary to non-overlapping parts of the Direct Repeat sequence so that when both primers hybridize to the same Direct Repeat and undergo elongation they will not be hindered by each other. In particular to avoid any hindrance during elongation reactions when one primer DRa is capable of elongation in the 5' Direction and the other primer DRb is capable of elongation in the 3' Direction the DRa is selected such that it is complementary to a sequence of the Direct Repeat located to the 5' side of the sequence of the Direct Repeat to which DRb is complementary. The primer used must have an oligonucleotide sequence capable of annealing to the consensus sequence of the Direct Repeat in a manner sufficient for amplification to occur under the circumstances of the particular amplification reaction. A person skilled in the art of amplification reactions will have no difficulty in determining which length and which degree of homology is required for good amplification

reactions to occur. The consensus sequence of the Direct Repeat of microorganisms belonging to the *M. tuberculosis* complex is given in sequence id. no. 2 and in figure 1.

In addition to what has already been disclosed in W095/31569 we also determined the spoligotypes of *M. tuberculosis* strains which were subcultured for many months both in the laboratory and in guinea pigs. The strains selected for this purpose were those used in a previous study on the stability of IS 6110 (2). All subcultured strains displayed the identical spoligotype patterns compared with the primary cultures thus indicating the pace of the molecular clock in this instance is slow enough for use in epidemiology of the disease.

Because of the large success and simplicity of the method for *Mycobacterium tuberculosis* strain differentiation and in view of problems in strain differentiation with other microorganisms we used the Direct Repeat consensus sequence to screen data bases with nucleic acid sequences from other microorganisms. Unfortunately no further matches were found. The Direct Repeat sequence appeared to be unique for the *Mycobacterium tuberculosis* as did their spacer sequences. As to date no function had actually been attributed to the Direct Repeat sequence it was unexpected that the sequence was universally distributed among other types of microorganisms. Such would at best be expected if the sequence had a function that was required also in other organisms.

Description of the invention

Notwithstanding the negative result after screening with the Direct Repeat consensus sequence we considered further analysis of known sequences by looking for a pattern in the nucleic acid sequences of other microorganisms reminiscent of the Direct Repeat-spacer pattern in *Mycobacterium tuberculosis*. Quite unexpectedly we found using a specifically designed computer programme that such patterns existed in a large number of other microorganisms with a broad range of genera. It appears that the DR-like sequences are very common in prokaryotes. They are however noticeably absent in eukaryotes. Chapter III of Bergeys Determinative Manual of Bacteriology Ninth edition (11) provides a table of characteristics for distinguishing prokaryotes from eukaryotes i.e. distinguish bacterium from microscopic eukaryotes in the shape of mold, yeast, algae or protozoans.

All bacterial sequences analysed revealed the presence of such

a sequence structure and thus the oligotyping method illustrated for *Mycobacterium tuberculosis* can be applied for differentiating between all strains of bacteria. It was totally unexpected that a consensus structure of this type could be universally found. The Direct Repeat sequences themselves are different between different genera but the general framework of a cluster of Direct Repeat sequences, separated by a number of non repetitive spacers is universally present in bacterial genomes. Considering the fact that thusfar no function has been attributed to such a region in *Mycobacterium tuberculosis* or in fact for any of the sequences comprising Direct Repeat like regions in any other bacteria for which such sequences had been described this is remarkable.

Bacteria can be divided into Archaeobacteria and Eubacteria. The eubacteria in turn can be distinguished into Gram-negative and Gram-positive bacteria with cell walls and Eubacteria lacking cell walls. Chapter IV of Bergeys determinative Manual of Bacteriology Ninth edition (11) reveals the characteristics for each group. Over a wide range of the subgroups in these 4 groups we have found the presence of the consensus structure i.e. the presence of DR-like loci. The IV groups have been subdivided by Bergey into more than 30 subgroups. We have examples in Groups 3,4,5 and 6, Group 11, 17, 31, 32, 33. The method according to the invention is particularly of interest for the bacteria that are pathogenic for humans. Group 4 comprises Gram negative bacteria. Genera from Group 4 are *Legionella* (which can cause pneumonia) and *Legionnaires* disease, the genus *Neisseria* (of which *Neisseria meningitidis* is well known as causative agent of meningitis and of which *Neisseria gonorrhoeae* is another example), the genus *Pseudomonas* (renown for hospital infections) and the genus *Bordetella* (of which *Bordetella pertussis* is well known as causative agent of whooping cough). In Group 5 bacteria as defined in Bergeys Manual the Enterobacteriaceae form a family of 30 genera. These bacteria form a particularly interesting group of Gram negative bacteria that infect humans. Suitable examples of genera from this family are *Enterobacter*, *Escherichia*, *Shigella*, *Salmonella*, *Serratia*, *Klebsiella* and *Yersinia*. Other less well known pathogenic Enterobacteriaceae genera are *Cedeca*, *Citrobacter*, *Kluyvera*, *Leclercia*, *Pantoea*, *Proteus*, *Providencia* and *Hafnia*. Other Group 5 families are Pasteurellaceae with the genus *Haemophilus* and the family Vibrionaceae with the genus *Vibrio*. *Haemophilus influenzae* is a leading cause of meningitis in children and also other septicemia conditions. *Vibrio cholerae* is the causative agent of cholera, *V. parahaemolyticus* can cause

food poisoning and *V. vulnificus* causes highly fatal septicemia.

Of the Enterobacteriaceae *Shigella*, *Escherichia* and *Salmonella* are best known and difficult to differentiate. *Shigella* is an intestinal pathogen of humans causing bacillary dysentery. Well known strains are *S. dysenteriae*, *S. flexneri*, *S. boydii*, *S. sonnei*. The genus *Salmonella* is well known for food poisoning. Well known *Salmonella* strains are *S. typhimurium*, *S. arizona*, *S. choleraesuis*, *S. bongori*. *Salmonella* are also causative agents of typhoid fever, enteric fevers, gastroenteritis and septicemia. The genus *Serratia* bacteria are opportunistic pathogens for hospitalized humans causing septicemia and urinary tract infections. Examples are *S. liquefaciens* and *S. marcescens*. Of the *Escherichia* *E. coli* is best known as major cause of urinary tract infections and nosocomial infections including septicemia and meningitis. Other species are usually associated with wound infections.

Enterobacter constitutes a problem genus of opportunistic pathogens causing burn wound and urinary tract infections occasionally also meningitis and septicemia. Well known species are *E. cloacae*, *E. sakazakii*, *E. aerogenes*, *E. agglomerans*, *E. gergoviae*. *Klebsiella* are also causative agents of bacteremia, pneumonia, urinary tract and other human infections in urological, neonatal, intensive care and geriatric patients. *Klebsiella pneumoniae* and *K. oxytoca* are examples of species in the genus.

Particularly interesting from a clinical point of view are also the Gram positive pathogenic bacteria. The genera *Streptococcus* and *Staphylococcus* form examples of such bacteria. *Streptococcus pneumoniae*, *Streptococcus pyogenes* and *Staphylococcus aureus* are examples thereof. Of the mentioned groups and genera the pathogenic bacteria are of interest. These bacteria are dangerous when infecting hospitals in particular.

Due to the increasing incidence of infection differentiation of potentially epidemiological organisms is also of interest. Such organisms comprise *Bordetella pertussis* and *Neisseria meningitidis* the causative organism of meningitis is of particular interest. Quite specifically pathogenic bacteria infecting hospitals and bacteria capable of causing epidemics are targets for the differentiation method according to the invention.

The invention consists of a method of in vitro amplification of nucleic acid using amplification primers in a manner known per se, in amplification reactions such as PCR, LCR or NASBA, wherein a pair of primers is used comprising oligonucleotide sequences sufficiently

complementary to a part of the Direct Repeat sequence of a bacterium other than a microorganism belonging to the *M. tuberculosis* complex of microorganisms for hybridisation to a Direct Repeat to occur and subsequently elongation of the hybridised primer to take place, said

5 primers being such that elongation in the amplification reaction occurs for one primer in the 5' Direction and for the other primer in the 3' Direction, wherein the Direct Repeat is a sequence with a length between 20-50 base pairs which occurs 5-60 times in a contiguous region of the bacterial genome, whereby the Direct Repeat sequences are separated by

10 spacer sequences with a length of between 20-50 nucleotides, said spacer sequences being non repetitive. By using the programme Patscan e.g. on the nucleic acid data bases for microorganism genomic sequences such motifs and thus also the identities of the various species specific Direct Repeats and the corresponding spacer sequences can be obtained. In

15 the Patscan programme the Direct Repeat can be designated p1 with a length between 20-50 basepairs then search for p1 20-50 basepairs downstream of p1. Thus this pattern in Patscan is described as $p1=(20..50)(20..50)p1(20..50)p1$. The length of the sequences can be varied as can the intermediate distance and the number of times the

20 Direct Repeat has to occur. A Direct Repeat can often have a length of 30-40 base pairs with a spacer length of 35-45 base pairs. Basically we looked for a stretch of identical repeat sequences interspersed by spacer sequences which do not necessarily share much of their sequence with the Direct Repeat of *M. tuberculosis*. The patscan programme is freely

25 accessible at the Internet site: <http://www-c.mcs.anl.gov/home/overbeek/-PatScan/HTML/patscan.html>. The programme was written by Ross Overbeek Mathematics and Computer Science Division Argonne National Laboratory Building 221 Room D-236 9700 S. Cass Avenue Argonne IL 60439 USA.

Most of the Repeats exhibit one or more of the following

30 characteristics, they end with a sequence similar to GAAAC i.e. exhibit at least 3 of the nucleotides of this consensus sequence at the terminus, preferably 4 or 5, start with CTTTG, have stretches of 3-4 identical bases. The termini can for example be selected from GAAAC, GAAXXC, GAACTC, GXAAC, GCAAC, GAAA, GAAXC, GAAGC and AAAC. Suitable Termini are provided

35 in Table II.

Organisms as diverse as the Archaeobacteria e.g. *Methanococcus jannasschi* (Group 31), *Haloferax mediterranei* (Group 33), the cyanobacteria *Calotrix* (Group 11), and *Anabeana* (Group 11), and purple bacteria e.g. *E.coli* (Group 5), *Mycobacterium tuberculosis* (Group 21) and

Thermus thermophilus (Group 4), Archaeoglobus (Group 32) and Thermotoga (Group 6) were found to possess DR-like sequences upon analysis of their genomes using the Patscan programme. In the subsequent study of literature from which these data were derived it also became clear from Southern blots that the Repeat sequences were also found in related species.

With regard to the genetic organisation the structures of the DR-like loci in the microorganisms is rather variable (figure 3). In *M. tuberculosis* the DR locus is large and in most isolates it is disrupted by an insertion element. This is also the case in *T. thermophilus*, however here the number of DVR's is only 11 and the DR locus is disrupted by two insertion elements. In *E. coli* K12 2 DR loci are present separated by approximately 22kb; in *Anabaena* the locus is of intermediate size and interrupted by a 130 bp sequence of unknown function or origin. In *H. mediterranei* the DR locus is of intermediate size and not disrupted, however there is evidence for a second DR locus on one of the mega plasmids found in this organism. In *M. jannaschii* there is one locus of intermediate size but at several other positions in the genome one or a few other DVR's are found. In most cases the DVR's are linked to a so-called Long Repeat (LR) element of unknown function. Also in *M. jannaschii* mega plasmids are found but in contrast to *H. mediterranei* they do not contain DR sequences.

Accession numbers for the sequences of various organisms for which the DR like loci have been found are provided here. For *E. coli* and *Shigella* M27059, M27060, U29579, U29580 and M18270. The relevant portions of the sequences are also disclosed by Blattner for *E. coli*. Nakata et al reveal in the Journal of Bacteriology (13) that downstream of the iap region a sequence of 29 bases appears 14 times 32 or 33 base pairs apart. Nucleotide sequences hybridizing to the 29 base pair sequence were also detected in *Shigella dysenteriae* and *Salmonella typhimurium*.

A DR-like sequence was found in the contig 214 of *S. pyogenes* M1(ATCC 700294) of the genome sequencing project of the University of Oklahoma. Further research into this DR-like sequence in other *S. pyogenes* revealed spacer polymorphism. The DR regions of eight *S. pyogenes* isolates were studied. The DR regions were isolated by PCR using primers that were derived from the database (University of Oklahoma, serotype M1 ATCC 700294). The sequence data is available under <http://www.genome.ou.edu>. This strain contains seven repeats and six spacers.

Five of the isolates gave a PCR product, these were a M2 strain, a M4 strain and three M1 strains. The M4 strain contained only a single repeat sequence that was flanked by the same sequences as the ATCC 700294. The M2 strain sequencing did not work, but the size of the PCR fragment indicated that two repeats are present. The three M1 strains were all the same, they contained four repeats and three spacers. The repeats were identical to ATCC 700294, while one of the spacers was identical to ATCC 700294 and two were different.

These studies on *S. pyogenes* show that the DR regions have conserved spacers and repeat sequences.

The Salmonella genomic sequence as sequenced by the University of Washington St Louis has also revealed the presence of DR-like sequences. The DR exhibits high homology with the Direct Repeat of *E. coli*. One of the contigs revealed 7 Repeats and 6 spacers.

A panel of five *E. coli* isolates and three *Shigella* strains were studied. The five *E. coli* isolates were selected to have an optimal diversity, they were isolated from different species or geographic regions. The *Shigella* strains are considered separate (sub)species. See Table 1. The isolates were obtained from the collection of Dr. Wim Gaastra.

Table 1

species	description	DRI*	DRII*
<i>E. coli</i> 184	American isolate	Southern	PCR
358	human urinary tract	Southern	Southern
968	mastitis	Southern	PCR
1008	chicken	PCR	PCR
1732	human intestine	Southern	PCR
<i>Shigella dysenteriae</i> 593		Southern	PCR
sonnei 595		Southern	PCR
boydii 603		Southern	PCR

* The DR regions were identified by Southern blot of genomic DNA and DRI and DRII regions of *E. coli* K12. When PCR is indicated the DR regions were identified by the Southern and the PCR. This PCR was done with primers derived from the K12 sequence.

The DRI and DRII sequences that could be amplified by PCR were

cloned and sequenced. Somehow the DRI regions could not be amplified by PCR using the primers designed on the K12 sequence, while the Southern data demonstrate that DRI is present. Apparently, the recognitions sites for the primers are polymorphic. The sizes of the DRII regions were found to vary greatly between these isolates. The smallest was a single repeat in the *S. sonnei* strain and the largest was a repeat cluster of at least 15 repeats in *E. coli* isolate 1008. The sequences of the repeats were highly conserved between these isolates. The *S. typhimurium* data is obtainable from the Internet <http://genome.wustl.edu/gsc/-bacterial/salmonella.html>.

The spacer sequences almost all were unique. Approximately 40 spacers have been sequenced and only three of them were already known from a previously sequenced DR region. This indicates a high number of different spacer sequences in *E. coli*.

Accession number X73453 provides the *Halorifax mediterranei* sequence. The sequence can also be found in Molecular Microbiology 17 of 1995 in an article by Mojica et al. (17). The Repeat sequence has also been found in related species.

The genomic project of the *Methanococcus jannaschii* reveals a DR-like sequence as is apparent from the Bult et al article in Science 273 of 1996 (18). The Accession number is U67459 i.a.

Accession number X87270 for *Anabeana sp* reveals 17 spacers and a LTRR element. These elements also occur in related species of cyanobacteria such as *Calotrix*. The sequence data are provided by Masepohl et al in BBA 1307 1996 (23).

Accession number AE000782 for *Archaeoglobus fulgidus* reveals three DR-like Repeats with the same Repeat sequence and the this has a slightly larger but closely related Repeat. The Repeats are present 20-30 times. The spacers are unique sequences. H.P. Klenk discloses sequence data in Nature 390 1997 (24).

The invention also covers a method of detection of a bacterium, said bacterium not belonging to the *M. tuberculosis* complex of microorganisms said method comprising

- 1) amplifying nucleic acid from a sample with the amplification method according to any of the preceding described embodiments of the amplification method according to the invention, followed by
- 2) carrying out a hybridisation test in a manner known per se, wherein the amplification product is hybridised to an oligonucleotide probe or a plurality of different oligonucleotide probes, each

oligonucleotide being sufficiently homologous to a part of a spacer of the Direct Region of the bacterium to be determined for hybridisation to occur to amplified product if such spacer nucleic acid was present in the sample prior to amplification, said hybridisation step optionally being carried out without prior electrophoresis or separation of the amplified product.

3) detecting any hybridised products in a manner known per se.

The method can be carried out in a manner such that the hybridisation test is carried out using a number of oligonucleotide probes, said number comprising at least a number of oligonucleotides probes specific for the total spectrum of bacteria it is desired to detect. In a suitable embodiment of a method according to the invention the oligonucleotide probe is at least seven oligonucleotides long and is a sequence complementary to a sequence selected from any of the spacer sequences of the Direct Repeat region of the bacterium to be determined or is a sequence complementary to fragments or derivatives of said spacer sequences, said oligonucleotide probe being capable of hybridising to such a spacer sequence and comprising at least seven consecutive nucleotides homologous to such a spacer sequence and/or exhibiting at least 60% homology, preferably exhibiting at least 80% homology with such a spacer sequence.

Preferably the method according to the invention is carried out to determine the presence and nature of a pathogenic bacterium selected from the group of Gram negative bacteria of Groups 4 and 5 of Bergeys Determinative Manual of Bacteriology ninth edition. Of particular interest due to damage caused by such pathogens are bacteria belonging to the families Enterobacteriaceae, Pasteurellaceae and Vibrionaceae of Group 5, most specifically the Enterobacteriaceae. Also of interest are the Gram positive bacteria of Group 17. Suitable examples of genera of the pathogenic bacterium to be detected from the group of Gram negative bacteria of Bergeys Determinative Manual of Bacteriology ninth edition are *Escherichia*, *Shigella*, *Salmonella*, *Klebsiella*, *Enterobacter*, *Yersinia*, *Serratia*, *Haemophilus*, *Vibrio*, *Legionella*, *Neisseria*, *Pseudomonas* and *Bordetella*. For the group of Gram positive bacterial genera *Staphylococcus* and *Streptococcus* are targets for the differentiation method.

Suitably in a method according to the invention for differentiating the type of bacterium in a sample, said bacterium not belonging to the *M. tuberculosis* complex the hybridisation pattern is

5 compared with that obtained with a reference. Such a reference can be the hybridisation pattern obtained with one or more known strains of the bacterium to be determined in analogous manner as the strain to be determined. Alternatively the reference is a source providing a list of spacer sequences and sources thereof, such as a data bank. Table II exhibits some suitable examples of sequences that occur as Direct Repeat sequences according to the invention for the genera illustrated.

Table II

Species	Repeat sequence	Reference	EMBL/Genbank accession number
<i>Mycobacterium tuberculosis</i>	GTCGTCAGACCCAAACCCCGAGAGGGGACGGAAAC	Hermans et al.	M27059 and M27060
<i>Escherichia coli</i>	CGGTTTATCCCGCTGGCGGGGAATC	Nakata et al. J. Bact. 171 3553-3556 (1989)	
<i>Shigella dysenteriae</i>	CGGTTTATCCCGCTGGCGGGGAATC	our own data	
<i>Shigella sonnei</i>	CGGTTTATCCCGCTGGCGGGGAATC	our own data	
<i>Shigella boydii</i>	CGGTTTATCCCGCTGGCGGGGAATC	our own data	
<i>Salmonella enteritidis</i>	CGGTTTATCCCGCTGGCGGGGAATC	our own data	
<i>Serratia marcescens</i>	CGGTTTATCCCGCTGGCGGGGAATC	our own data	
<i>Salmonella typhimurium</i>	CGGTTTATCCCGCTGGCGGGGATACAC	contig 70A06 of the typhimurium genome project. Univ. of Washington St. Louis	Contig 214 of the S. pyogenes genome project University of Oklahoma
<i>Streptococcus pyogenes</i>	GTTTATAGAGCTATGCTGTTTGAATGGTCCCAAACT		

(Table II)

Species	Repeat sequence	Reference	EMBL/Genbank accession number
<i>Thermus aquaticus</i> <i>thermophilus</i> <i>Thermotoga maritima</i>	AATCCCCCTTACGGGGCTCAATCCCTTGCAA GTTTCAATACTTCCTTAGAGGTATGGAAC	Ashby et al. Plasmid 24 1-11 (1990) Liebl. et al. Microbiol. 142 2533- 2542 (1996)	M33159 Z69341
<i>Anabaena</i>	GTTTAACTAACAATAATCCCTATCAGGGATTGAAAC	Masepohl BBA 1307 26- 30 1996	X87270
<i>Calotrix</i> <i>Haloferax mediterranei</i>	GTTTAAACTTTATAAAATCCCTTTTAGGGATTGAAAC GTTACAGACGAAACCTTAGTGGTTGAAGC	idem Mojica et al. Molec. Microbiol. 17 85-93 (1995)	Z47161 X7453
<i>Methanococcus jannaschii</i>	AATTAAATCAGACCGTTTCGGAATGGAAA	Bult et al. Science 273 1058-1073 (1996)	U67459 (for the large DR-like region)
<i>Methanobacterium</i> <i>thermoautotrophicum</i> <i>Archaeoglobus fulgidus</i>	ATTTCATCCCATTTTGGTCTGATTTTAAC GTTAAATCAGACCAAAATGGGATTGAAAT CTTTCATCCCATTTTGGTCTGATTTCAAC	Smith et al. J. Bact. 179 7135-7155 (1997) Klenk et al. Nature 390 364-370 (1997)	AE000920 (for R2) AE000878 (for R1) AE000782 (whole genome)

Not only the above methods fall within the scope of the invention but also specifically selected primer pairs for carrying out such a method. A pair of primers according to the invention is a pair wherein both primers comprise oligonucleotide sequences of at least 7
5 oligonucleotides and are sufficiently complementary to a part of the Direct Repeat sequence of the microorganism *E. coli* for hybridisation to occur and subsequently elongation of the hybridised primer to take place, said primers being such that elongation in the amplification reaction occurs for one primer in the 5' Direction and for the other primer in the
10 3' Direction and wherein sufficiently complementary means said oligonucleotide sequence comprises at least seven consecutive nucleotides homologous to such a Direct Repeat sequence and/or exhibits at least 60% homology, preferably at least 80% homology, most preferably more than 90% homology with the corresponding part of the Direct Repeat sequence.
15 Suitable Direct Repeat sequences are provided in Table II. In particular such a primer pair can comprise one primer DRa capable of elongation in the 5' Direction and the other primer DRb capable of elongation in the 3' Direction with DRa being complementary to a sequence of the Direct Repeat located to the 5' side of the sequence of the Direct Repeat to which DRb
20 is complementary, the Direct Repeat being present in the Direct Region of *E. coli*. Another suitable pair comprises primers with oligonucleotide sequences of at least 7 oligonucleotides and are sufficiently complementary to a part of the Direct Repeat sequence of the microorganism *S. typhimurium* for hybridisation to occur and subsequently
25 elongation of the hybridised primer to take place, said primers being such that elongation in the amplification reaction occurs for one primer in the 5' Direction and for the other primer in the 3' Direction and wherein sufficiently complementary means said oligonucleotide sequence comprises at least seven consecutive nucleotides homologous to such a
30 Direct Repeat sequence in particular the Sequence provided in Table II and/or exhibits at least 60% homology, preferably at least 80% homology, most preferably more than 90% homology with the corresponding part of the Direct Repeat sequence. In particular such a pair comprises one primer DRa capable of elongation in the 5' Direction and the other primer DRb
35 capable of elongation in the 3' Direction with DRa being complementary to a sequence of the Direct Repeat located to the 5' side of the sequence of the Direct Repeat to which DRb is complementary, the Direct Repeat being present in the Direct Region of *S. typhimurium*.

Kits for carrying out a differentiation method according to any

of the described embodiments also fall within the scope of the invention. Such kits comprise a primer pair according to any of the described embodiments and optionally an oligonucleotide probe or a carrier, said carrier comprising at least 1 oligonucleotide probe specific for a spacer region of a bacterium to be determined said bacterium not belonging to *M tuberculosis* complex, preferably the oligonucleotide probe as defined, being an oligonucleotide probe of at least 10 nucleotides, preferably more than 12 nucleotides, in particular comprising between 12 to 40 nucleotides, said probe being sufficiently homologous to any of the spacer sequences or to fragments or derivatives of such spacer sequences to hybridise to such a spacer sequence, said oligonucleotide probe comprising at least 10 consecutive nucleotides homologous to such a spacer sequence and/or exhibiting at least 60% homology, preferably exhibiting at least 80% homology, most preferably exhibiting more than 90% homology with the corresponding part of the spacer sequence. Suitably a kit according to the invention comprises a data carrier with required reference patterns of the bacterial strain to be determined.

REFERENCES

1. Beck-Sagué, C., S.W. Dooley, M. D. Hutton, J. Otten, A. Breeden, J.T. Crawford, A.E. Pitchenik, C. Woodley, G. Cauthen, W.R. Jarvis. 1992 Hospital outbreak of multidrug-resistant *Mycobacterium tuberculosis* infections. JAMA. 268: 1280-1286.
2. Van Soolingen, D., P.W.M. Hermans, P.E.W. de Haas, D. R. Soll, and J.D.A. van Embden. 1991. The occurrence and stability of insertion sequences in *Mycobacterium tuberculosis* complex strains; evaluation of IS-dependent DNA polymorphism as a tool in the epidemiology of tuberculosis. J. Clin. Microbiol. 29:2578-2586.
3. CDC. 1992. Transmission of multidrug-resistant tuberculosis among immunocompromised persons in a correctional system. New York 1991. MMWR 41: 507-509.
4. CDC. 1991. Nosocomial transmission of multidrug resistant tuberculosis among HIV-infected persons, Florida and New York, 1988-1991. MMWR 40: 585-591.
5. Cave, M.D., K.D. Eisenach, P.F. McDermott, J.H. Bates, and J.T.

- Crawford. 1991. IS6110: Conservation of sequence in the *Mycobacterium tuberculosis* complex and its utilization in DNA fingerprinting. *Mol. Cell. Probes* 5:73-80.
- 5 6. Daley, C.L. P.M. Small, G.F. Schecter, G.K. Schoolnik, R.A. McAdam, W.R. Jacobs, Jr., and P.C. Hopewell. 1992. An outbreak of tuberculosis with accelerated progression among persons infected with the human immunodeficiency virus: an analysis using restriction fragment length polymorphisms. *N. Engl. J. Med.* 326:231-235.
 - 10 7. Dooley, S.W., M.E. Villarino, M. Lawrence, et al. 1992. Nosocomial transmission of tuberculosis in a hospital unit for HIV-infected patients. *JAMA.* 267:2632-2634.
 - 15 8. Dwyer, B., K. Jackson, K. Raios, A. Sievers, E. Wilshire and B. Ross. 1993. DNA Restriction fragment analysis to define an extended cluster of tuberculosis in homeless men and their associates. *J. Inf. Diseases,* 167: 490-494.
 - 20 9. Edlin, B.R., J.I. Tokars, M.H. Grieco, J.T. Crawford, J. Williams, E.M. Sordillo, K.R. Ong, J.O. Kilburn, S.W.. Dooley, K.G. Castro, W.R. Jarvis and S.D. Holmberg. 1992. An outbreak of multidrug-resistant tuberculosis among hospitalized patients with the acquired immunodeficiency syndrome. *N. Eng. J. Med.* 326:1514-1521.
 - 25 10. Fischl, M.A, R.B. Uttamchandani, G.L. Daikos, et al. 1992. An outbreak of tuberculosis caused by multiple-drug resistant tubercle bacilli among patients with HIV infection. *Ann. Intern. Med.* 117:177-183.
 - 30 11. Bergeys Determinative Manual of Bacteriology Ninth Edition
 12. Groenen, P.M.A., A.E. van Bunschoten, D. van Soolingen, and J.D.A. van Embden. 1993. Nature of DNA polymorphism in the direct repeat cluster of *Mycobacterium tuberculosis*; Application for strain differentiation by a novel method. *Mol. Microbiol.* (1993) 10 (5) 1057-1065.
 - 35 13. Nakata et al *Journal of Bacteriology* June 1989 p3553-3556.
 14. Hermans, P.W.M., D. van Soolingen, J.W. Dale, A.R. Schuitema, R.A. McAdam, D. Catty, and J.D.A. van Embden. 1990. Insertion element IS986 from *Mycobacterium tuberculosis*: a useful tool for diagnosis and epidemiology of tuberculosis. *J. Clin. Microbiol.* 28:2051-2058.
 15. Hermans, P.W.M., D. van Soolingen, E.M. Bik, P.E.W. de Haas,

- J.W. Dale, and J.D.A. van Embden. 1991. The insertion element IS987 from *M. bovis* BCG is located in a hot spot integration region for insertion elements in *M. tuberculosis* complex strains. *Infect. Immun.* 59:2695-2705.
- 5 16. Hermans, P.W.M., D. van Soolingen, and J.D.A. van Embden. 1992. Characterization of a major polymorphic tandem repeat in *Mycobacterium tuberculosis* and its potential use in the epidemiology of *Mycobacterium kansasii* and *Mycobacterium gordonae*. *J. Bacteriol.* 174:4157-4165.
- 10 17. Mojica et al., *Molecular Microbiology* 17, 1995
18. Bult et al., *Science* 273, 1996
19. Mazurek, G.H., M.D. Cave, K.D. Eisenach, R.J. Wallace JR, J.H. Bates, and J.T. Crawford. 1991. Chromosomal DNA fingerprint patterns produced with IS6110 as strain specific markers for epidemiologic study of tuberculosis. *J. Clin. Microbiol.* 29:2030-2033.
- 15 20. McAdam, R.A., P.W.M. Hermans, D. van Soolingen, Z.F. Zainuddin, D. Catty, J.D.A. van Embden, and J.W. Dale. 1990. Characterization of a *Mycobacterium tuberculosis* insertion sequence belonging to the IS3 family. *Mol. Microbiol.* 4:1607-1613.
- 20 21. Mendiola, M.V., C. Martin, I. Otal, and B. Gicquel. 1992. Analysis of regions responsible for IS6110 RFLP in a single *Mycobacterium tuberculosis* strain. *Res. Microbiol.* 143:767-772.
- 25 22. Otal, I., C. Martin, V. Vincent-Lévy-Frébault, D. Thierry, and B. Gicquel. 1991. Restriction fragment length polymorphism analysis using IS6110 as an epidemiological marker in tuberculosis. *J. Clin. Microbiol.* 29:1252-1254.
23. Masepohl et al., *kBBA* 1307 1996
- 30 24. Klenk, H.P., *Nature* 390, 1997
25. Palittapongarapim, P.S., S. Rienthong, and W. Panbangred. 1993. Comparison of restriction fragment length polymorphism of *M. tuberculosis* isolated from cerebrospinal fluid and sputum: a preliminary report. *Tubercle and lung disease* 1993. 74, 204-207.
- 35 26. Ross, C., K. Raios, K. Jackson, and B. Dwyer. 1992. Molecular cloning of a highly repeated element from *Mycobacterium tuberculosis* and its use as an epidemiological tool. *J. Clin. Microbiol.* 30:942-946.

27. Small, P.M., R.W. Schafer, P.C. Hopewell, S.P. Singh, M.J. Murphy, E. Desmond, M.F. Sierra, and G.K. Schoolnik. 1993. Exogenous reinfection with multidrug-resistant *M. tuberculosis* in patients with advanced HIV infection. *N. Eng. J. Med.* 328:1137-1144.
28. Thierry, D., M.D. Cave, K.D. Eisenach, J.T. Crawford, J.H. Bates, B. Gecquel, and J.L. Guesdon. 1990. IS610, an IS-like element of *M. tuberculosis* complex. *Nucleic Acids Res.* 18:188.
29. Van Embden, J.D.A., D. van Soolingen, P.M. Small, and P.W.M. Hermans. 1992. Genetic markers for the epidemiology of tuberculosis. *Res. Microbiol.* 143: 385-391.
30. Van Embden, J.D.A., M.D. Cave, J.T. Crawford, J.W. Dale, K.D. Eisenach, B. Gicquel, P.W.M. Hermans, C. Martin, R. McAdam, T.M. Shinnick, and P.M. Small. 1993. Strain identification of *Mycobacterium tuberculosis* by DNA fingerprinting; Recommendations for a standardized Methodology *J. Clin. Microbiol.* 31:406-409.
31. Van Soolingen, D., P.W.M. Hermans, P.E.W. de Haas, and J.D.A. van Embden. 1992. Insertion element IS1081-associated Restriction Fragment Length Polymorphism in *Mycobacterium tuberculosis* Complex species: a reliable tool for recognizing *Mycobacterium bovis* BCG. *J. Clin. Microbiol.* 30:1772-1777.
32. Van Soolingen, D., P.E.W. de Haas, P.W.M. Hermans, P.M.A. Groenen and J.D.A. van Embden. 1993. Comparison of various repetitive DNA elements as genetic markers for strain differentiation and epidemiology of *Mycobacterium tuberculosis*. *J. Clin. Microbiol.* 31:1987-1995.
33. Yuen L.K., B.C. Ross, K.M. Jackson, and B. Dwyer. 1992. Characterization of *Mycobacterium tuberculosis* strains from Vietnamese patients by southern blot hybridization. *J. Clin. Micro.* 31: 1615-1618.
34. Kamerbeek J., L.M. Schouls, A. Kolk, M. van Agterveld, D. van Soolingen, S. Kuijper, J.E. Bunschoten, H. Molhuizen, R. Shaw, M. Goyal and J.D.A. van Embden 1997. Simultaneous detection and strain differentiation of *Mycobacterium tuberculosis* for diagnosis and epidemiology. *J. Clin. Microbiol.* 35: 907-914.

DESCRIPTION OF THE FIGURES

- Figure 1 depicts the structure of the DR region of *M. bovis* BCG as determined previously by Hermans et al. and Groenen et al. (12, 15). For the sake of convenience we will designate a DR plus its 3'adjacent spacer sequence as a "Direct Variant Repeat" (DVR). Thus, the DR region is composed of a discrete number of DVR's, each consisting of a constant part (DR) and a variable part (the spacer).
- Figure 2 depicts multiple, short synthetic oligomeric DNA sequences based on the sequences of the unique spacer DNA's within the DR region.
- Figure 3 shows the genetic organisation of the structures of the DR-like loci in various bacterial species.
- depicts the transcription direction of open reading frame (ORF)
- For *M. tuberculosis*: MTCY 16B7.26, 27 and 30C are unknown genes/proteins.
- For *E. coli*: iap gene function is alkaline phosphatase isozyme conversion. ORF f94, f305, YGCE and f223 are unknown genes/proteins.
- For *S. pyogenes*: ORF1 and 2 are unknown genes/proteins.
- For *T. thermophilus*: ORFC and D are unknown genes/proteins and ORF 1A and 1B are possibly transposases of IS elements 1000 and 1000A.
- For *Anabaena*: No ORFs were annotated in the flanking sequences. The 130 bp insert is of unknown origin.
- For *Haloferax mediterranei*: ORF21 is an unknown gene/protein. Probably another repeat cluster is also present on the megaplasmid pHM500.
- For *Methanococcus jannaschii*: Comprises about 10 repeat clusters, the largest one of which comprises 25 repeats. All repeat clusters are coupled to a Long Repeat (LR) segment of 425bp. There are 18 LR's, some of which contain only one repeat. Smaller LR segments are also present, ΔLR. In one case, a cluster contains 5 repeats without LR (see ref. 18)
- For *M. thermoautotrophicum*: Two repeat clusters SRI and SRII flanked by

LRI, LRII. LRI and LRII are almost identical and are homologues of the LR segment of *M. jannaschii*. SRI and SRII are separated by 500 kb in the genome.

- 5 For *Thermatoga maritima*: CelA gene encodes cellulase: endo-1,4-beta-glucanase (EC 3.2.1.4) and CelB is also a cellulase exhibiting 58% identity with celA.

- 10 For *Archaeoglobus fulgidus*: The SRIA and SRIB repeat clusters have the same Repeat Sequence and the SRII Repeat Sequence is also clearly homologous. The SR clusters are separated by about 400bp. SRIB and SRII are located near tRNA genes. SRIA lies adjacent to an unknown ORF3.

Figure 4

- 15 Hybridization Patterns of 17 *E. coli* isolates. Thirty four different spacer oligonucleotides were covalently linked to a membrane and PCR amplified DNA of *E. coli* was hybridized as described (Kamerbeek et al. 1997), except that the primers used to amplify the DR locus were specific for the DR sequence from *E. coli*. Note the polymorphism observed in *E.*
20 *coli* due to the strain-dependent presence or absence of spacer DNA.

Figure 5

- Hybridization Patterns of 4 *Salmonella typhimurium* isolates. Six
25 different spacer oligonucleotides were covalently linked to a membrane and PCR amplified *Salmonella* DNA was hybridized as described (Kamerbeek et al 1997), except that the primers used to amplify the DR locus were specific for the DR locus of *E. coli*. Note the polymorphism observed in *Salmonella* due to the strain-dependent presence or absence of spacer DNA.

30

09/647596

PCT/NL98/00186


WO 99/51771

1/8

Fig 1

M. bovis BCG

IS6110

*M. tuberculosis* H37Rv

Consensus DR sequence: GTGTCAGACCCAAACCCCGAGAGGGACGGAAC


09/647596

WO 99/51771


PCT/NL98/00186

2/8

Fig 2



amplification products :



09/647596


WO 99/51771

PCT/NL98/00186

Fig 3.1


M. TUBERCULOSIS

H37Rv




ESCHERICHIA COLI

K12



STREPTOCOCCUS PYOGENES



3/8


WO 99/51771

PCT/NL98/00186

Fig 3.2


THERMUS AQUATICUS THERMOPHILUS

HB8 ATCC 27634




ANABAENA sp. PCC.7120

4/8



HALOFERAX MEDITERRANEI



WO 99/51771


09/647596

PCT/NL98/00186


5/8

Fig 3.3

METHANOCOCCUS JANNASCHII



METHANOBACTERIUM THERMOAUTOTROPHICUM




WO 99/51771


6/8

Fig 3₄

THERMOTOGA MARITIMA



ARCHAEOGLOBUS FULGIDUS



WO 99/51771

09/647596

PCT/NL98/00186

7/8

Fig 4




Fig. 6

SEQ ID No. 1: GTCGTCAGACCCAAAACCCCGAGAGGGGACGGAAAC

SEQ ID No. 2: CGGTTTATCCCCGCTGGCGCGGGAACTC

SEQ ID No. 3: CGGTTTATCCCCGCTGGCGCGGGGAACTC

SEQ ID No. 4: CGGTTTATCCCCGCTGGCGCGGGGAACTC

SEQ ID No. 5: CGGTTTATCCCCGCTGGCGCGGGGAACTC

SEQ ID No. 6: CGGTTTATCCCCGCTGGCGCGGGGAACTC

SEQ ID No. 7: CGGTTTATCCCCGCTGGCGCGGGGAACTC

SEQ ID No. 8: CGGTTTATCCCCGCTGGCGCGGATACAC

SEQ ID No. 9: GTTTTAGAGCTATGCTGTTTTGAATGGTCCCAAAC

SEQ ID No. 10: AATCCCCTTACGGGGCTCAATCCCTTGCAA

SEQ ID No. 11: GTTTCATACTTCCTTAGAGGTATGGAAAC

SEQ ID No. 12: GTTTTAACTAACAAAATCCCTATCAGGGATTGAAAC

SEQ ID No. 13: GTTTAAACTTTATAAAATCCCTTTTAGGGATTGAAAC

SEQ ID No. 14: GTTACAGACGAACCCTAGTTGGGTGGAAGC

SEQ ID No. 15: AATTAAAATCAGACCGTTTCGGAATGGAAA

SEQ ID No. 16: ATTTCAATCCCATTTTGGTCTGATTTTAAC

SEQ ID No. 17: GTTAAAATCAGACCAAATGGGATTGAAAT


SEQ ID No. 18: CTTTCAATCCCATTTTGGTCTGATTTCAAC

WO 99/51771

09/647596
PCT/NL98/00186

8/8

Fig 5



09/647596

H434

Fig. 6

SEQ ID No. 1: GTCGTCAGACCCAAAACCCCGAGAGGGGACGGAAAC

SEQ ID No. 2: CGGTTTATCCCCGCTGGCGCGGGGA⁶ACTC

SEQ ID No. 3: CGGTTTATCCCCGCTGGCGCGGGGA⁶ACTC

SEQ ID No. 4: CGGTTTATCCCCGCTGGCGCGGGGA⁶ACTC

SEQ ID No. 5: CGGTTTATCCCCGCTGGCGCGGGGA⁶ACTC

SEQ ID No. 6: CGGTTTATCCCCGCTGGCGCGGGGA⁶ACTC

SEQ ID No. 7: CGGTTTATCCCCGCTGGCGCGGGGA⁶ACTC

SEQ ID No. 8: CGGTTTATCCCCGCTGGCGCGGATACAC

SEQ ID No. 9: GTTTTAGAGCTATGCTGTTTGAATGGTCCCAA⁶ACT

SEQ ID No. 10: AATCCCCTTACGGGGCTCAATCCCTTGCAA

SEQ ID No. 11: GTTCAATACTTCCTTAGAGGTATGGAAAC

SEQ ID No. 12: GTTTTAACTAACAAAAATCCCTATCAGGGATTGAAAC

SEQ ID No. 13: GTTTAACTTTATAAAATCCCTTTTAGGGATTGAAAC

SEQ ID No. 14: GTTACAGACGAACCCTAGTTGGGTTGAAGC

SEQ ID No. 15: AATTAAAAATCAGACCGTTTCGGAATGGAAA

SEQ ID No. 16: ATTTCAATCCCATT¹⁷TTGGTCTGATTTTAAC

~~SEQ ID No. 17:~~ GTTAAATCAGACCAA¹⁷AATGGGATTGAAAT

SEQ ID No. ~~18~~¹⁷: CTTTCAATCCCATT¹⁷TTGGTCTGATTTCAAC

AMENDED SHEET

COMBINED DECLARATION AND POWER OF ATTORNEY

(ORIGINAL DESIGN, NATIONAL STAGE OF PCT OR CIP APPLICATION)

As a below named inventor, I hereby declare that

My residence, post office address and citizenship are as stated below next to my name, I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled:

A method of interstrain differentiation of bacteria

the specification of which: (complete (a), (b) or (c) for type of application)

REGULAR OR DESIGN APPLICATION

- a. ☐ is attached hereto.
b. ☐ was filed on _____ as Application
Serial No. _____ and was amended on _____
(if applicable)

PCT FILED APPLICATION ENTERING NATIONAL STAGE

- c. ☒ was described and claimed in International application No. PCT/NL98/00186
filed on 3 April 1998
and as amended on _____ (if any)

ACKNOWLEDGEMENT OF REVIEW OF PAPERS AND DUTY OF CANDOR

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, paragraph 1.56(a).

In compliance with this duty there is attached an information
disclosure statement 37 CFR 1.97

PRIORITY CLAIM

I hereby claim foreign priority benefits under Title 35, United States Code paragraph 119 of any foreign application (s) for patent of inventor's certificate listed below and have also identified below any foreign application for patent of inventor's certificate having a filing date before that of the application on which priority is claimed.

(complete (d) or (e))

- d. ☐ no such applications have been filed
 e. ☐ such applications have been filed as follows

**EARLIEST FOREIGN APPLICATION(S), IF ANY FILED WITHIN 12 MONTHS
 (6 MONTHS FOR DESIGN) PRIOR TO SAID APPLICATION**

Country	Application Number	Date of filing (day, month, year)	Date of Issue (day, month, year)	Priority claimed

**ALL FOREIGN APPLICATION(S), IF ANY FILED MORE THAN 12 MONTHS
 (6 MONTHS FOR DESIGN) PRIOR TO SAID APPLICATION**

CONTINUATION-IN-PART

(Complete this part only if this is a continuation-in-part application)

I hereby declare claim the benefit under Title 35, United States code, paragraph 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claim of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, paragraph 112, I acknowledge the duty to disclose material information as defined in Title 37, Code of Federal Regulations, paragraph 1.56(a) which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

(Application Serial No.)	(Filing date)	(Status)	(patented, pending, abandoned)
(Application Serial No.)	(Filing date)	(Status)	(patented, pending, abandoned)


POWER OF ATTORNEY

As a named inventor, I hereby appoint the following attorney(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith: Robert J. PATCH, Reg. No. 17,355, Andrew J. PATCH, Reg. No. 32,925, Robert F. HARGEST, Reg. No. 25,590, Benoit CASTEL, Reg. No. 35,041, Eric Jensen, Reg. No. 37,855, and Thomas W. PERKINS, Reg. No. 33,027 c/o YOUNG & THOMPSON, Second Floor, 745 South 23rd Street, Arlington, Virginia 22202.

Address all telephone calls to Young & Thompson at 703/521-2297.


I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that wilful false statements and the like so made are punishable by fine or imprisonment, or both under Section 1001 of Title 18 of the United States Code and that such wilful false statements may jeopardize the validity of the application or any patent issued thereon.

1-00
Full name of sole or first inventor: VAN EMBDEN, Johannes Dirk Anthonie
Inventor's signature




Date 13 November 2000 Country of Citizenship: The Netherlands
Residence: The Netherlands
Post Office Address: Van Limburg Stirumstraat 15, NL-3581 VB UTRECHT NLX

2-00
Full name of second inventor: SCHOOLS, Leendert Marinus
Inventor's signature



Date 13 November 2000 Country of Citizenship: The Netherlands
Residence: The Netherlands
Post Office Address: IJsselsteen 47, NL-3961 GB WIJK BIJ DUURSTEDE NLX

3-00
Full name of third inventor: JANSEN, Rudolph
Inventor's signature



Date 21-november 2000 Country of Citizenship: The Netherlands
Residence: The Netherlands
Post Office Address: Golfpark 149, NL-8241 AC LELYSTAD NLX

CHECK PROPER BOX(ES) FOR ANY ADDED PAGE(S) FORMING A PART OF THIS DECLARATION